

MLP -09

Age-Related Changes in Serum Biochemical Parameters in the Domestic Rabbit (*Oryctolagus cuniculus*)

K.A. Adebisi

Department of Animal Science, University of Ibadan, Ibadan

E-mail: karamatsanusi@yahoo.co.uk

Abstract

The influence of age on serum biochemical components in the domestic rabbit was investigated. The serum total protein (g/dL), albumin (g/dL), globulin (g/dL), creatinine (mg/dL), urea mg/dL, cholesterol (mg/dL), Alkaline phosphatase (ALP; iu/l), Aspartate aminotransferase (AST; iu/l) and Alanine aminotransferase (ALT; iu/l) were assessed in 24 male rabbits at 7, 14 and 21 weeks of age to represent weaners, growers and pubertal stages. Data were analysed using analysis of variance at $\alpha_{0.05}$ and descriptive statistics. Results of serum biochemistry revealed a steady significant increase in total protein (4.2 ± 0.3 ; 5.3 ± 0.5 ; 6.9 ± 0.2), albumin (2.2 ± 0.7 ; 2.8 ± 0.4 ; 4.0 ± 0.2), globulin (1.9 ± 0.3 ; 2.5 ± 0.4 ; 2.9 ± 0.2), cholesterol (49.5 ± 1.1 ; 56.4 ± 4.2 ; 62.4 ± 5.4) and creatinine (0.8 ± 0.1 ; 1.2 ± 0.1 ; 1.9 ± 0.3) at 7, 14 and 21 weeks respectively. Serum urea levels were similar across the ages. The values recorded for ALP and AST at 7 weeks (39.6 ± 7.5 ; 101.1 ± 6.7) and 14 weeks (38.7 ± 6.4 ; 104.5 ± 8.7) were significantly higher than the 21 weeks (33.5 ± 2.7 ; 95.8 ; 5.9) while ALT at 7 weeks (27.33 ± 5.6) was significantly lower than 14 (38.2 ± 3.9) and 21 weeks (37.4 ± 3.0). The range of values recorded at 7 weeks for total protein, albumin and globulin were lower than the ranges reported in literature while the range recorded for the other parameters were within that reported in the literature. The age of the rabbits had a significant effect on their serum biochemical parameters.

Keywords: Rabbit, age, serum biochemical values, serum reference values

Introduction

The domestic rabbit is used extensively in medical and biological research because they are mammals, small in size and have relatively short gestation period and generation interval. They can also be kept for meat, wool and as a pet. Rabbits have received a considerable research attention in Nigeria. This stems from the current trend in the country to exploit some unpopular but potential meat resource to complement the conventional ones like cattle, sheep, poultry or pigs. These researches are usually aimed at how best to produce numerous cheap rabbits, within the shortest possible time at a relatively low cost without compromising health and performance. When such studies are carried out, usually, rabbits of any age, sex or physiological status may be used and the blood profile (haematology or serum biochemistry) is usually one of the tools used to monitor the study. This creates a need to establish reference values for the blood profile to serve as a guide in interpreting results from such research. In establishing such values, it is important to bear in mind that several factors such as analytical methods, age of animal, environmental factors or seasonal influence may affect results apart from the physiological status of the animal. These therefore, necessitate the establishment of reference values for our local use and consider how they vary with age. For technical reasons, many biochemical measurements are more conveniently made on serum rather than plasma but the concentrations of most analytes are effectively the same in both fluids (Marshall, 2000).

A lot of studies have been carried out on the blood profile of various domestic animals, but there is dearth of information on the reference haematological and serum biochemical values for rabbits in the warm humid environment (Chineke *et al.*, 2002) and more especially, considering the influence of age. Ozegbe *et al* (2000), Ozegbe *et al* (2000), Awojobi and Opiah (2002) have documented changes in plasma or serum profile at some physiological stages for sows, cows and doe rabbits respectively. Tambuwal *et al.* (2002) have reported some serum biochemical values for goats considering age and sex. For rabbits, Chineke *et al.* (2002) considered some breeds and crosses while Iwuji *et al.* (2017) also reported serum biochemical values for rabbits at different locations.

The objective of this study was therefore to assess the influence of age at an actively growing phase of male rabbits on some serum biochemical components.

Materials and Methods

The study was carried out at the Teaching and Research farm and the Physiology laboratory Department of Animal Science, University of Ibadan. A total of twenty-four New Zealand x Chinchilla

crossbred male rabbits were used for the study. They were purchased from a breeder farm at 5 weeks of age immediately after weaning. On arrival of the animals, they were given vitamins and necessary prophylaxis and allowed to equilibrate for about two weeks afterwards to provide time for a homeostatic return to physiological normality and to stabilize their nutritional and environmental conditions. Rabbits were checked regularly for signs of illness to ensure only healthy animals were used in the course of the study. They were housed individually in cages. Pelleted concentrate rations containing 19% crude protein, 10% crude fibre and 2,480Kcal digestible energy/Kg feed was provided ad-lib along with *Tridax procumbens* as forage. Fresh clean water was also supplied ad-lib. Blood samples for serum biochemistry were taken at intervals of seven weeks. The first was at 7 weeks of age, second at 14 weeks and third at 21 weeks representing the weaners, growers and pubertal stages respectively. Blood sample taken from each animal was analysed separately and taken as a replicate. The experimental design was a completely randomized design and analysis of variance was done to check for significant difference of each of the blood parameters across the ages.

The rabbits were deprived of feed for about twelve hours prior to collection of blood samples to minimize the effect of feed on the concentration of the various blood constituents (Mitruka and Rawnsley, 1977). Blood was sampled from the marginal vein of the ear. About 5mL of blood from individual rabbits was collected into separate labelled sterile tubes. Serum was separated about an hour after clotting by centrifuging for about 30 minutes. The serum was drawn off by pipette and immediately frozen until analysis was carried out. The serum constituents analysed for were total protein, albumin, globulin, Alkaline phosphatase (ALP), Aspartate aminotransferase (AST), Alanine aminotransferase (ALT), creatinine, urea and cholesterol. Total protein was determined by biuret reaction, albumin by bromocresol green reaction and globulin by subtracting albumin from total protein as described by Toro and Ackermann (1979). Urea was determined by the use of enzyme urease and cholesterol by direct serum method as described by Burtis and Ashwood (1996). The enzymes and creatinine were determined indirectly and the reactions monitored spectrophotometrically at fixed time intervals (Burtis and Ashwood, 1996).

The data obtained were subjected to analysis of variance of SAS (1999) and means were separated using Duncan multiple range test.

Results and Discussion

The mean serum biochemical values for the rabbits are presented in Table 1. The results indicated a consistent significant ($p < 0.05$) increase in mean values for total protein (4.18 ± 0.26 ; 5.32 ± 0.50 ; 6.88 ± 0.23), albumin (2.19 ± 0.72 ; 2.82 ± 0.36 ; 4.03 ± 0.24), globulin (1.86 ± 0.33 ; 2.50 ± 0.38 ; 2.86 ± 0.24), cholesterol (49.52 ± 1.13 ; 56.44 ± 4.24 ; 62.36 ± 5.41) and creatinine (0.79 ± 0.10 ; 1.23 ± 0.10 ; 1.87 ± 0.30) at ages 7, 14 and 21 weeks respectively. The increase in these values with respect to age is as with reference values for humans in which case values set at different age groups from new-born to adulthood for total protein, albumin, cholesterol and creatinine increase steadily with the increase in age (Burtis and Ashwood, 1996). Results on increase in protein also confirm reports from Chineke *et al.* (2002) in which serum total protein increased from a mean of 4.5 at 42 days to 6.8 at 91 days, albumin, 3.18 to 3.78 and globulin 1.23 to 2.42 respectively.

Table 1: Mean ($\pm$ SD) serum biochemical values of rabbits as influenced by age

Serum components	Age of Rabbits		
	7 weeks	14 weeks	21 weeks
Total protein (g/dL)	4.18 ± 0.26^c	5.32 ± 0.50^b	6.88 ± 0.23^a
Albumin (g/dL)	2.19 ± 0.72^c	2.82 ± 0.36^b	4.03 ± 0.24^a
Globulin (g/dL)	1.86 ± 0.33^c	2.50 ± 0.38^b	2.86 ± 0.24^a
Creatinine (mg/dL)	0.79 ± 0.10^c	1.23 ± 0.10^b	1.87 ± 0.30^a
Urea (mg/dL)	24.26 ± 2.51	26.65 ± 5.06	25.21 ± 4.72
Cholesterol (mg/dL)	49.52 ± 1.13^c	56.44 ± 4.24^b	62.36 ± 5.41^a
ALP (iu/L)	39.60 ± 7.54^a	38.70 ± 6.44^a	33.50 ± 2.70^b
AST (iu/L)	101.14 ± 6.66^a	104.48 ± 8.71^a	95.81 ± 5.94^b
ALT (iu/L)	27.33 ± 5.64^b	38.19 ± 3.91^a	37.39 ± 2.97^a

Means across the rows with different superscripts are significantly different ($p < 0.05$), SD- Standard Deviation, ALP – Alkaline phosphatase; AST- Aspartate aminotransferase; ALT- Alanine aminotransferase.

The increase in creatinine supports the fact that creatinine is expected to increase from infancy to puberty, parallel to the development of skeletal muscle as this is the period growth is most prominent. The

increase in cholesterol confirm reports from Marshal (2000) that blood cholesterol increase with age. Values recorded for urea were similar across the ages. It ranged from 24.26 ± 2.51 (7 weeks) to 26.65 ± 5.06 (14 weeks). This is probably because blood urea levels in health are more affected by diet than other factors. The values recorded for ALP and AST at 7 weeks (39.60 ± 7.54 ; 101.14 ± 6.66) and 14 weeks (38.70 ± 6.44 ; 104.48 ± 8.71) were significantly higher than 21 weeks (33.50 ± 2.70 ; 95.81 ± 5.94) while ALT at 7 weeks (27.33 ± 5.64) was significantly lower than 14 (38.19 ± 3.91) and 21 weeks (37.39 ± 2.97). The results of ALP, AST and ALT supports reports of Burtis and Ashwood (1996) that the serum activity of most serum enzymes decrease from childhood to puberty in humans but that of ALT may continue to rise at least in men until middle age.

The range of serum biochemical values recorded at the three stages of growth is presented in Table 2. The range recorded at 7 weeks for total protein, albumin and globulin were lower than the ranges reported in the literature for those parameters. This may partly be a result of the significantly lower mean values recorded at that stage in Table 1. This further stresses the fact that the parameters may be highly influenced by age and the need to classify these reference values according to age groups. The range recorded for the other parameters were within that reported in the literature.

Table 2: Range of serum biochemical values of rabbits recorded at different ages

Serum components	Age of rabbits			¹ Range in literature	² Range in literature
	7 weeks	14 weeks	21 weeks		
Total protein(g/dL)	3.86 – 4.58	4.50 – 5.75	6.58 – 7.22	5.00 - 8.00	5.40 – 7.30
Albumin (g/dL)	2.00 – 2.27	2.10 – 3.40	3.58 – 4.59	2.80 - 4.00	2.40 – 4.50
Globulin (g/dL)	1.42 – 2.44	1.94 – 3.09	2.27 – 3.29	2.20 - 4.00	2.90 – 4.90
Creatinine(mg/dL)	0.60 – 0.95	0.80 – 1.40	1.16 – 2.33	-	0.50 – 2.20
Urea (mg/dL)	19.30 – 28.90	20.60 – 36.60	17.50 – 32.80	9.00 - 32.00	10.00 – 33.00
Cholesterol(mg/dL)	47.50 – 51.30	50.30 – 62.00	56.00 – 75.00	20.00 - 83.00	10.00 – 80.00
ALP (iu/L)	31.80 – 50.10	33.00 – 51.00	32.00 – 40.20	10.00 - 90.00	4.00 – 20.00
AST (iu/L)	82.35 – 109.00	99.00 – 111.40	84.40 – 108.00	42.00 - 98.00	10.00 – 120.00
ALT (iu/L)	20.00 – 30.00	30.20 – 43.00	30.90 - 41.00	49.00 - 79.00	10.00 – 45.00

¹CCAC (1980), ²Rosenthal (2003), ALP – Alkaline phosphatase; AST- Aspartate aminotransferase; ALT- Alanine aminotransferase.

Conclusion and Recommendation

Age influences the serum biochemical components of rabbits and the most important influence that determines the overall effect of age are the degree of sexual maturity and the amount of skeletal muscle mass of the animal. Since significant differences were observed for virtually all serum biochemical components assessed in the study at different ages, it will be necessary to group rabbits according to age when setting reference values for them to avoid misguidance in interpreting laboratory results. Results from this study can therefore, serve as a contribution to reference serum biochemical values for rabbits of different age groups.

References

- Awojobi, H.A. and Opiah, G.O. (2002). The effect of physiological status on some blood parameters of the New Zealand white doe rabbits. *Proc. 7th ann. Conf. Anim. Sci. Ass. of Nig. (ASAN), Sept. 16-19, 2002, UNAAB, Nigeria.*
- Burtis, C.A. and Ashwood, E.R. (1996). *Tietz fundamental of clinical chemistry*, 4th edition. W. B. Saunders Company, Philadelphia, Pennsylvania 19106.
- CCAC(1980). Guide to the care and use of experimental animals. *Canadian Council on Animal Care*, 1: 82-90.
- Chineke, C.A., Adeniran, F.A., Ologun, A.G., Ikeobi, C.O.N. and Oseni, O.A.(2002). Analysis of some serum biochemical parameters in New Zealand white rabbits and their crosses. *Proc. 27th Ann. Conf. Nig. Soc. Of Anim. Prod. (NSAP), March 17 – 21, 2002, Fed. Univ. of Tech. Akure. Nig. Pp: 5 – 7.*
- Iwuji, T.C., Nwapi, D.O., Ogbuwu I.P., Kadurumba, E.O., Egenuka, F.C. and Okere, P.C. (2017). Management, blood and reproductive parameters of rabbits reared in Imo State. *Nig. J. Anim. Prod.*, 44(2); 230-247.
- Marshall, W.J.(2000). *Clinical chemistry*, 4th edition. Harcourt Publishers Ltd, Harcourt place, London.
- Mitruka, B.M. and Rawnsley, H.M.(1977). Clinical biochemical and hematological reference values in normal experimental animals. MASSON publishing, USA, Inc.

- Ozegbe P.C., Michael, A.N. and Oke, O.A. (2000). Biochemical changes in the plasma of crossbred sows. *Trop. J. Anim. Sci.*,3(2); 35-41.
- Ozegbe, P.C., Ajadi, R.A. and Ajeneye, V.O. (2000). Influence of lactation on some plasma biochemical parameters in N'dama and white Fulani cows. *Trop. J. Anim. Sci.*,3(2): 43-49.
- Rosenthal, K. (2002). Rabbit reference ranges (blood chemistry). University of Pennsylvania School of Veterinary Medicine. Downloaded online; www.cal.vet.upenn.edu
- SAS, (1999). Statistical Analysis System user's guide. Statistical Analysis System Institute Inc., Cary, North Carolina 27513, USA
- Tambuwal, F.M., Ajagbe, B.M. and Bangana, B. (2002). Hematological and biochemical values of apparently healthy Red Sokoto goats. *Proc. 27th Ann. Conf. Nig. Soc. Of Anim. Prod. (NSAP)*, March 17-21, 2002, Fed. Univ. Akure, Nigeria.
- Toro, G. and Akermann, P.G. (1975). *Practical clinical chemistry*, 1st edition. Little Brown and Co. Boston, USA.