

UTILIZATION OF BITTER LEAF (*VERNONIA AMYGDALINA*) MEAL AS FEED FOR WEST AFRICAN DWARF (WAD) GOATS

A. O. OLOSUNDE AND S. M. ODEYINKA

Department of Animal Sciences, Obafemi Awolowo University, Ile-Ife.

*Correspondence: mrbgodveng@gmail.com, 08032371575

ABSTRACT

This study determined the utilization and the optimum level of offer of bitter leaf (*Vernonia amygdalina*) meal (BLM) to West African Dwarf (WAD) goats fed graded levels of bitter leaf meal. In a 20-week trial, twenty four West African Dwarf (WAD) goats of both sexes, 5-7 months old, were randomly allotted to four treatments of graded levels of BLM (0% BLM (Control diet), 15% BLM, 30% BLM and 45% BLM diets) in a completely randomized design to determine the utilization of bitter leaf meal as feed for West African Dwarf goats. The diets served as supplements to a basal ration of *Panicum maximum*. Two digestibility trials were carried out. Data obtained were statistically analyzed with the General Linear Model and significant means were separated using the Duncan's Multiple Range Test. The digestible ether extract intake and digestible organic matter of goats on 0% BLM were significantly ($P<0.05$) higher than the values obtained for goats on bitter leaf meal diets. The average daily weight gain (g/day) for goats on 0% BLM was significantly ($P<0.05$) higher than that of others. There were significant ($P<0.05$) differences in the percentage mortality of the goats fed experimental diets. The percentage mortality of goats on 30% and 45% BLM were significantly ($P<0.05$) higher than that of others. It was concluded that bitter leaf meal can be included in the diets of (WAD) goats up to 15% without any deleterious effect.

Keywords: Bitter leaf meal, Utilization, *Panicum maximum*, West African Dwarf Goats.

INTRODUCTION

One of the major problems confronting the small ruminant production is the non-availability of feed all year round to meet the maintenance and productive requirements of the animals. Although, grasses abound in the tropics, seasonal changes which affect their palatability and nutritive value had been a major problem in ruminant animal production (Alokan, 1998). Poor productivity and high mortality of stock, which characterize this industry is largely explained by the inadequacy of feeding the right quantity and quality of feeds to the various livestock species (Odeyinka, 2014).

The unprecedented increase in the cost of conventional ingredients (e.g. maize) used in compounding livestock feeds has necessitated intensive investigations into the use of agricultural and agro-industrial by-products, leguminous fodder trees and shrubs which are regarded as non-conventional feed sources (Mousa, 2011). Bitter leaf (*Vernonia amygdalina*) is a drought tolerant shrub with petiole leaf of about 6mm diameter. The leaves are used for human consumption, suggesting that the excess may be utilized as feed (Daodu and Babayemi, 2009). Bitter leaf is a

manageable size shrub that is available during dry season, its nutritional, economic value and potential as a dry season feed source for ruminant animals is underutilized. Hence, this study was designed to determine the utilization of bitter leaf meal as a feedstuff for West African Dwarf goats.

MATERIALS AND METHODS

The experiment was carried out at the Sheep and Goat Unit of the Teaching and Research Farm, Obafemi Awolowo University, Ile-Ife, Osun State. Twenty four West African Dwarf (WAD) weaner goats of both sexes were used in this experiment. The ages of the animals were between 5 and 7 months. The goats were quarantined dewormed with levamisole and ivomec injection prior to the commencement of the experiment. The animals were randomly allotted to four treatments in a completely randomized design with six goats per treatment and provided fresh feed and water daily. The experiment lasted for a period of 20 weeks. The *Vernonia amygdalina* leaves for this study were harvested from the cut branches of the planted shrub within the environment of the Teaching and Research farm, Obafemi

Awolowo University, Ile-Ife. Samples were air dried for two days and sun dried on concrete slabs for a day. The samples were ground prior to its incorporation in goat diets. Four concentrate diets comprising 0, 15, 30 and 45% of Bitter Leaf (*Vernonia amygdalina*) Meals (BLM) were compounded and fed to the goats as supplements to a basal ration of guinea grass (*Panicum maximum*) (Table 1). The goats were fed based on 3% of their body weights. The feed was offered once daily at 09.00 hour. Feed refusals were collected and weighed the following morning before offering fresh feed. This was used to estimate feed intakes. The animals were subjected to 14 day adaptation period. The animals were weighed at the beginning of the experiment and weekly thereafter. This was to assess live weight changes. Two digestibility trials were carried out at (10th week) and (16th week) of the experimental period. The metabolism trial was conducted with 3 goats per treatment in metabolism cages with facilities for separate collection of faeces and urine. The proximate composition of the feed and fecal sample was determined by standard methods (AOAC, 1990). Data obtained were statistically analyzed with the General Linear Model of SAS (2008) and the Duncan New Multiple Range Test option of SAS (2008) was used to detect significant differences among means.

RESULTS AND DISCUSSION

The results of the chemical composition of experimental diets are presented in Table 2. The crude protein and crude fibre content of the diets increased with increase in the inclusion level of BLM while decrease in ether extract and nitrogen free extract values were observed. The crude protein levels of all the diets were above the 9.89% minimum requirement for ruminants reported by Ogunbosoye and Babayemi, 2010. The result showed that bitter leaf meal (BLM) can serve as suitable source of nutrients in livestock feed. The results of apparent digestibility coefficient of the dry matter and nutrients intake of experimental diets are shown in Table 3. There was significant difference in the dry matter digestibility ($P<0.05$) among the experimental animals. The dry matter digestibility, (71.53%) from the CD diet was significantly ($P<0.05$) higher than mean values

of 56.89 and 58.97%, obtained by feeding 15% BLM and 30% BLM respectively. The apparent digestibility coefficient of the crude protein, crude fibre, ether extract, nitrogen free extract and ash were significantly ($P<0.05$) affected by the inclusion levels of bitter leaf meal. The crude protein digestibility results obtained in this study was similar to the report of Ngi *et al.* (2006) who obtained a range of 54.01 – 64.49% CP digestibility for Maradi × West African Dwarf cross bred goats. The performance assessment of the goats on the test diets showed that goats on control diet (CD) had higher dry matter, crude protein, ether extract and organic matter digestibilities. This could be attributed to the high fiber content of the diets with bitter leaf inclusion. Table 4 showed the mean dry matter and nutrient intake (g/day) of goats fed the experimental diets for a 20-week period. The mean daily dry matter intakes (DMI) (g/day) were not significantly ($P>0.05$) different. The average daily intake (g/day) of the animals were 297.10, 282.93, 276.94 and 256.03g for animals fed 30% BLM, 45% BLM, CD and 15% BLM respectively. The values (256.03-297.10 DMI g/day) obtained in this study were lower than 305.20 - 311.50 g/day reported by Odedire and Oloidi (2014). The mean live weight gains (g/day/animal), from animals fed CD was significantly ($P<0.05$) higher than animals fed 30% BLM, 15% BLM and 45% BLM, respectively. The average daily weight gains were 13.57, 9.88, 5.66 and 3.75 g/day/animal for goats fed CD, 30% BLM, 15% BLM and 45% BLM diets, respectively (Table 4). This strengthens the belief that weight reduction was indeed a consequence of the ingested *Vernonia amygdalina* leaves. Atangwho *et al.* (2012) reported respective decrease of 12.78% and 38.51% in body weight gain, of *V. amygdalina* fed rats at 5% and 15% against 17.45% decrease in body weight gain of orlistat at end of study ($P<0.05$); but with no effect on feed intake at 4-week. The weight gains from this study were higher than 1.47-2.67 g/day reported by Osakwe and Udeogu (2007). There were significant ($P<0.05$) differences in the FCR of the goats across the diets. The results revealed the ability of goats on control diet to convert the feed consumed to weight gain. Isah *et al.*

(2013) reported similar FCR range of 31.89-62.56 for West African Dwarf goats fed different tropical browse plants with *Pennisetum purpureum* as basal diet. There were significant ($P<0.05$) differences in the percentage mortality of the goats fed experimental diets. The percentage mortality of goats on 30% and 45% BLM were significantly ($P<0.05$) higher than that of others.

CONCLUSION

It could be concluded that 15% bitter leaf meal inclusion is the optimum level that should be offered in the diet of West African Dwarf goats for maximal performance without any detrimental effect on the animals.

REFERENCES

- Alokan, J. A. 1998. Intakes and digestibility by Yankasa sheep of diets containing different proportions of legumes and grass (*Cynodon nlemfuensis*). In: O. O. Oduguwa, A. O. Fanimu and O. A. Osinowo (eds). *Proceedings of Silver Anniversary of Nigerian Society for Animal Production*. pp: 337-338.
- AOAC 1990. Official Method of Analysis. 15th Edition, Association of Official Analytical Chemists, Washington D.C., USA.
- Atangwho, I. J., Edet, E. E., Uti, D. E., Obi, A. U., Asmawi, M. Z. and Ahmad, M. 2012. Biochemical and histological impact of *Vernonia amygdalina* supplemented diet in obese rats. *Saudi Journal of Biological Sciences*, 19(3): 385-392.
- Daodu, M. O. and Babayemi, O. J. 2009. Utilization of Some Edge-Row Plants as Forage in Nigeria. *Pakistan Journal of Nutrition*, 8 (8): 1269-1274.
- Isah, O. A., Oguntayo, S. A., Dawodu, R. O., Diya, O. O., Afolabi, M. O. and Omoniyi, L. A. 2013. Feed Utilization, Rumen Parameters and Microbial Profile of Goats fed Different Tropical Browse Plants with *Pennisetum purpureum* as Basal Diet. *The Pacific Journal of Science and Technology*. Vol. 14(1): 297-405.
- Mousa, M. R. H. 2011. Effect of Feeding *Acacia* as Supplements on the Nutrient Digestion, Growth Performance, Carcass Traits and Some Blood Constituents of Awassi Lambs under the Conditions of North Sinai. *Asian Journal of Animal Sciences*, 5: 102-117.
- Ngi, J., Ayoade, J. A. and Oluremi, O. I. A. 2006. Evaluation of dried cassava leaf meal and maize offal as supplements for goats fed rice straw in dry season. *Livestock Research For Rural Development*. Volume 18, Article 127. <http://www.lrrd.org/lrrd18/9/ngi18127.htm>
- Odedire, J. A. and Oloidi, F. F. 2014. Feeding Wild Sunflower (*Tithonia Diversifolia* Hemsl., A Gray) to West African Dwarf Goats as a Dry Season Forage Supplement. *World Journal of Agricultural Research* 2.6 (2014): 280-284.
- Odeyinka, S. M. 2014. That they may eat all year round: The intellectual odyssey of a ruminant nutritionist. An inaugural lecture delivered at Oduduwa Hall, Obafemi Awolowo University, Ile-Ife, Nigeria on Tuesday 14th January, 2014. pp 7-25.
- Ogunbosoye, D. O. and Babayemi, O. J. 2010. Potential values of some non-leguminous browse plants as dry season feed for ruminants in Nigeria. *African Journal of Biotechnology* Vol. 9 (18), pp. 2720-2726.
- Osakwe, I. I. and Udeogu, R. N. 2007. Feed intake and nutrient digestibility of West African Dwarf (WAD) goat fed *Pennisetum purpureum* supplemented with *Gmelina arborea*. *Animal Research International* (2007) 4(3): 724-727.
- SAS 2008. *SAS User's Guide Statistics*, SAS Inc. Cary., North Carolina, 2008 edition.

Table 1: Composition of Experimental Diets

Ingredients	0% BLM	15% BLM	30% BLM	45% BLM
Corn bran	45.00	30.00	15.00	-
Bitter leaf meal	-	15.00	30.00	45.00
Brewers dried grain	30.00	30.00	30.00	30.00
Palm kernel cake	22.50	22.50	22.50	22.50
Bone Meal	1.50	1.50	1.50	1.50
Salt	0.50	0.50	0.50	0.50
Vitamin Premix	0.50	0.50	0.50	0.50
Total	100.00	100.00	100.00	100.00
Calculated % Crude protein	14.70	17.14	19.58	22.01

Table 2: Chemical composition of diets fed to experimental goats

Parameter	0% BLM	15% BLM	30% BLM	45% BLM
Dry matter (DM)	85.70	90.18	87.71	91.04
Analysis % of DM				
Organic matter	88.19	90.60	90.11	86.67
Crude protein	20.92	23.53	25.44	29.33
Crude fibre	8.38	9.67	10.10	10.68
Ether extract	10.16	8.96	7.20	5.61
Ash	11.81	9.40	9.89	13.33
Nitrogen free extract	43.62	43.68	42.27	41.07

Table 3: Apparent digestibility coefficient of the dry matter and nutrient intake (g/day) of goats fed experimental diets

Parameter %	0% BLM	15% BLM	30% BLM	45% BLM	SEM	PROB
DDMI	71.53 ^a	56.89 ^b	58.97 ^b	64.71 ^{ab}	1.71	0.004
DCPI	73.57 ^a	60.68 ^b	59.80 ^b	68.44 ^a	1.69	0.003
DCFI	77.53 ^b	57.18 ^c	63.87 ^c	85.67 ^a	2.57	<0.0001
DEEI	82.91 ^a	68.89 ^c	63.19 ^d	74.55 ^b	1.75	<0.0001
DNFE	93.82 ^b	94.22 ^b	96.78 ^a	79.94 ^c	1.40	<0.0001
DASH	57.23 ^a	40.49 ^b	41.66 ^b	61.81 ^a	2.39	<0.0001
DOM	85.06 ^a	76.02 ^b	77.18 ^b	77.90 ^b	1.00	0.001

^{a,b,c}: Means within each row with different superscript are significantly different ($p \leq 0.05$)

SEM: Standard error of mean; PROB: Probability; DDMI: Digestible dry matter intake; CPI: Digestible crude protein intake; DCFI: Digestible crude fibre intake; DEEI: Digestible ether extract intake; DNFE:

Digestible nitrogen free extract intake; DASH: Digestible ash; DOM: Digestible organic matter

Table 4: Performance characteristics of the experimental goats

Parameter	CD	15% BLM	30% BLM	45% BLM	SEM	PROB
Average daily feed consumption (g/day)						
Concentrate	139.12	124.75	145.89	138.38	5.15	0.56
Panicum	137.74	131.63	151.21	144.54	5.52	0.65
Total	276.86	256.38	297.10	282.93	10.24	0.59
Average initial live weight (kg)	8.28	8.38	8.10	8.05	0.27	0.97
Average final weight (kg)	10.47	9.18	9.90	8.71	0.40	0.44
Mean live weight (kg)	9.52	8.78	9.21	8.48	0.27	0.55
Total weight gain (kg)	1.90 ^a	0.79 ^c	1.38 ^b	0.53 ^d	0.13	<0.0001
Average daily gain (g)	13.57 ^a	5.66 ^c	9.88 ^b	3.75 ^d	0.94	<0.0001
Percentage mortality (%)	0.00 ^c	0.00 ^c	50.00 ^a	33.33 ^b	4.41	<0.0001
Feed conversion ratio (FCR)	20.40 ^d	45.34 ^b	30.07 ^c	75.45 ^a	4.48	<0.0001

^{a,b,c,d}: Means within each row with different superscript are significantly different ($p \leq 0.05$)

SEM: Standard error of mean; PROB = Probability